Author Index

Abbott, F.S., see Chan, G.L.-Y. 295
Abdalla, T., see Greene, S.V. 121
Abou-Khalil, S.
—, Abou-Khalil, W.H., Masoud, A.N. and Yunis, A.A.
High-performance liquid chromatographic determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and ultraviolet detection 111
Abou-Khalil, W.H., see Abou-Khalil, S. 111
Andersson, T., see Kågedal, B. 261
Axelson, J.E., see Chan, G.L.-Y. 295
Baars, A.M., see Vree, T.B. 458
Badarani, M.M., see Dubruc, C. 208
Balmér, K.
—, Zhang, Y., Lagerström, P.-O. and Persson, B.-A.
Determination of metoprolol and two major metabolites in plasma and urine by column liquid chromatography and fluorometric detection 357
Bargar, E.M.
Application of column switching in high-performance liquid chromatographic analysis of medroxalol in plasma 143
Bauer, J.H., see Souther, M.E. 27
Bayne, W.F., see Rothwell, C.M. 447
Bekersky, I., see Puglisi, C.V. 347
Belanger, K.
—, Collins, J.M. and Klecker, Jr., R.W.
Technique for detection of DNA nucleobases by reversed-phase high-performance liquid chromatography optimized for quantitative determination of thymidine substitution by iododeoxyuridine 57
Benecke, R.
—, Vetter, B. and De Zeeuw, R.A.
Rapid micromethod for the analysis of mitotane and its metabolite in plasma by gas chromatography with electron-capture detection 287
Benowitz, N.L., see Jacob, III, P. 277
Bhamra, R.K.
—, Ward, A.E. and Holt, D.W.
Column liquid chromatographic measurement of betaxolol in plasma or serum 229
Brinkman, U.A.Th., see Nielen, M.W.F. 159
Bryan, C.S., see Greene, S.V. 121
Cantelmi, M.G., see Palmerini, C.A. 378
Carlsson, M., see Kågedal, B. 261
Chan, G.L.-Y.
—, Axelson, J.E., Abbott, F.S., Kerr, C.R. and McErlane, K.M.
Determination of propafenone in biological fluids by fused-silica capillary gas chromatography using electron-capture detection 295
Chervet, J.P., see Humbert, H. 319
Collins, J.M., see Belanger, K. 57
Conte, Jr., J.E.
— and Zurlinden, E.
Column liquid chromatographic determination of cefpiramide in human serum and urine 452
Curini, R., see Laganá, A. 135
Dann, O., see Gluth, W.P. 331
D'Ascenzo, G., see Laganá, A. 135
Davis, P.C.
Determination of xamoterol in human plasma by high-performance liquid chromatography with electrochemical detection 233
De Grood, P.M.R.M., see Vree, T.B. 458
Denneberg, T., see Kågedal, B. 261
Denouel, J., see Humbert, H. 319
De Vries, J.X.
—, Walter-Sack, I. and Ittensohn, A.
Analysis of benzbromarone in human plasma and urine by high-performance liquid chromatography and gas chromatography–mass spectrometry 420
De Zeeuw, R.A., see Benecke, R. 287
Dollery, C.T., see Richmond, R. 241
Dubruc, C.
—, Hermann, Ph., Haddouche, A., Badarani, M.M. and Thénot, J.J.
Determination of S-carboxymethyl-L-cysteine in plasma by high-performance liquid chromatography with column switching following precolumn derivatization 208
Elias-Baker, B.A., see Jacob, III, P. 277
Ema, M., see Itami, T. 168
Falardeau, P., see Martineau, A. 1
Farjam, A., see Nielen, M.W.F. 159
Farrant, M.
—, Zia-Gharib, F. and Webster, R.A.
Automated pre-column derivatization with *o*-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase liquid chromatography 385

Feely, J., see O'Reilly, T. 190
Ferrara, F.J., see Puglisi, C.V. 347
Fini, C., see Palmerini, C.A. 378
Floridi, A., see Palmerini, C.A. 378
Fouda, H.G.
—, Ronfeld, R.A. and Weidler, D.J.
Gas chromatographic–mass spectrometric analysis and preliminary human pharmacokinetics of sertraline, a new antidepressant drug 197
Frei, R.W., see Nielen, M.W.F. 159
Frydman, A., see Le Liboux, A. 151
Fujiwara, K., see Kobayashi, K. 439
Gaillot, J., see Le Liboux, A. 151
Gehrke, C.W., see Souther, M.E. 27
Gibson, D.M., see Wright, D.S. 223
Gluth, W.P.
— and Dann, O.
Determination of fluorescent diamidines in plasma of experimental animals by high-performance liquid chromatography 331
Grech-Belanger, O.
—, Leboeuf, E. and Langlois, S.
Assay of diltiazem and deacetyldiltiazem by capillary gas chromatography 89
Greene, S.V.
—, Abdalla, T., Morgan, S.L. and Bryan, C.S.
High-performance liquid chromatographic analysis of vancomycin in plasma, bone, atrial appendage tissue and pericardial fluid 121
Grood, P.M.R.M. de, see Vree, T.B. 458
Guthrie, R., see Ho, J. 269
Haddouche, A., see Dubruc, C. 208
Hal, H.J.M. van, see Paanakker, J.E. 203
Hamazaki, H., see Iwase, H. 403
Hatsumura, H., see Sayo, H. 129
Hatta, M., see Hirano, T. 371
Heavey, D., see Richmond, R. 241
Heilmann, P., see Schöneshöfer, M. 434
Hermann, Ph., see Dubruc, C. 208
Hirai, K., see Kominami, G. 216
Hirano, T.
—, Hatta, M., Oka, K. and Wakasugi, K.
Liquid chromatographic determination of serum lipid peroxides as diphenylthiobarbituric acid-reactive substance to collagen disease patients – a preliminary study 371
Ho, J.
—, Guthrie, R. and Tieckelmann, H.
Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by high-performance liquid chromatography with fluorimetric detection 269
Holt, D.W., see Bhamra, R.K. 229
Hoppe, A., see Kågedal, B. 261
Hosokawa, M., see Sayo, H. 129
Hotta, K., see Iwase, H. 403
Hsieh, J.Y.-K., see Rothwell, C.M. 447
Humbert, H.
—, Denouel, J., Chervet, J.P., Lavene, D. and Kiechel, J.R.
Determination of sub-nanogram amounts of dihydroergotamine in plasma and urine using liquid chromatography and fluorimetric detection with off-line and on-line solid-phase drug enrichment 319
Ickrath, O., see Ress, K. 465
Imashuku, S., see Tsunamoto, K. 414
Ingen, H.E. van, see Nielen, M.W.F. 159
Ishii, I., see Iwase, H. 403
Itami, T.
—, Ema, M., Sakamoto, J. and Kawasaki, H.
Microheterogeneity of rat α-fetoprotein on a copper chelate column 168
Ito, T.
—, Matsuki, Y., Kurihara, H. and Nambara, T.
Sensitive method for determination of captopril in biological fluids by gas chromatography–mass spectrometry 79
Ittensohn, A., see De Vries, J.X. 420
Iwai, K., see Shibati, K. 173
Iwase, H.
—, Ishii, I., Hamazaki, H. and Hotta, K.
Determination of uridine 5′-diphosphate glucose–collagen glucosyltransferase activity by high-performance liquid chromatography and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 403
Jacob, III, P.
—, Elias-Baker, B.A., Jones, R.T. and Benowitz, N.L.
Determination of benzoylecgonine and cocaine in biologic fluids by automated gas chromatography 277
Jimenez, J., see Sonesson, A. 366
Johnson, C.A., see Vanscoik, K.G. 183
Jones, R.T., see Jacob, III, P. 277
Jordan, R.A., see Wright, D.S. 223
Kågedal, B.
—, Andersson, T., Carlsson, M., Denneberg, T. and Hoppe, A.
High-performance liquid chromatography of 2-mercaptopropionylglycine and its metabolite 2-mercaptopropionic acid in plasma and urine after treatment with thio-

pronine 261
Karlsson, L.
Direct injection of urine on a high-performance liquid chromatographic column-switching system for determination of Δ^9-tetrahydrocannabinol-11-oic acid with both ultraviolet and electrochemical detection 309
Kawada, T., see Shibati, K. 173
Kawasaki, H., see Itami, T. 168
Keeling, P.W.N., see O'Reilly, T. 190
Kerr, C.R., see Chan, Ç.L.-Y. 295
Kiechel, J.R., see Humbert, H. 319
Klecker, Jr., R.W., see Belanger, K. 57
Kobayashi, K.
—, Sakoguchi, T., Fujiwara, K., Taniuchi, K., Kohri, K. and Matsuoka, A.
High-performance liquid chromatographic determination of amphotericin B in human urine 439
Kohri, K., see Kobayashi, K. 439
Kominami, G.
—, Matsumoto, H., Nishimura, R., Yamaguchi, T., Konaka, R., Sugeno, K., Hirai, K. and Kono, M.
Combined high-performance liquid chromatography and enzyme immunoassay for active metabolites of a sleep inducer (rilmazafone), a ring-opened derivative of benzodiazepines, in human plasma 216
Konaka, R., see Kominami, G. 216
Kono, M., see Kominami, G. 216
Krämer, B., see Ress, K. 465
Krungkrai, J.
—, Yuthavong, Y. and Webster, H.K.
High-performance liquid chromatographic assay for pteroylpolyglutamate hydrolase 47
Kuo, K.C., see Souther, M.E. 27
Kurihara, H., see Ito, T. 79
Labrude, P., see Menu, P. 397
Laganá, A.
—, Curini, R., D'Ascenzo, G., Marino, A. and Rotatori, M.
High-performance liquid chromatographic determination of norfloxacin in human tissues and plasma with fluorescence detection 135
Lagerström, P.-O., see Balmér, K. 357
Langlois, S., see Grech-Belanger, O. 89
Larsson, L., see Sonesson, A. 11, 366
Lauff, J.J.
Ion-pair high-performance liquid chromatographic procedure for the quantitative analysis of theophylline in serum samples 99
Lavene, D., see Humbert, H. 319
Leboeuf, E., see Grech-Belanger, O. 89
Le Liboux, A.
—, Frydman, A. and Gaillot, J.
Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by reversed-phase high-performance liquid chromatography 151
Liboux, A. Le, see Le Liboux, A. 151
Liebich, H.M., see Ress, K. 465
Linari, F., see Mentasti, E. 253
Lumpkin, R.H., see Souther, M.E. 27
MacCrehan, W.A.
— and Schönberger, E.
Reversed-phase high-performance liquid chromatographic separation and electrochemical detection of retinol and its isomers 65
McErlane, K.M., see Chan, G.L.-Y. 295
Mac Mathúna, P., see O'Reilly, T. 190
Maessen, P.A.
—, Mross, K.B., Pinedo, H.M. and Van der Vijgh, W.J.F.
Improved method for the determination of 4′-epidoxorubicin and seven metabolites in plasma by high-performance liquid chromatography 339
Maltby, N., see Richmond, R. 241
Marangella, M., see Mentasti, E. 253
Marino, A., see Laganá, A. 135
Martineau, A.
— and Falardeau, P.
Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary gas chromatography–negative-ion chemical-ionization mass spectrometry 1
Masoud, A.N., see Abou-Khalil, S. 111
Mathúna, P. Mac, see O'Reilly, T. 190
Matsuki, Y., see Ito, T. 79
Matsumoto, H., see Kominami, G. 216
Matsumoto, S., see Togari, A. 41
Matsuoka, A., see Kobayashi, K. 439
Mentasti, E.
—, Savigliano, M., Marangella, M., Petrarulo, M. and Linari, F.
High-performance liquid chromatographic determination of glyoxylic acid and other carbonyl compounds in urine 253
Menu, P.
—, Pauly, F., Labrude, P. and Vigneron, C.

Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate–haemoglobin species by high-performance liquid chromatographic resolution 397
Michida, T., see Sayo, H. 129
Minelli, A., see Palmerini, C.A. 378
Morgan, S.L., see Greene, S.V, 121
Mross, K.B., see Maessen, P.A. 339
Nagatsu, T., see Togari, A. 41
Nakazawa, H.
Fluorimetric determination of rat growth hormone in pituitary cell culture by high-performance liquid chromatography with post-column derivatization 409
Nambara, T., see Ito, T. 79
Newton, C.J.
—, Roberts, N.B. and Taylor, W.H.
Separation of human pepsins in gastric juice by high-performance ion-exchange chromatography 391
Nielen, M.W.F.
—, Van Soest, R.E.J., Van Ingen, H.E., Farjam, A., Frei, R.W. and Brinkman, U.A.Th.
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by liquid chromatography with precolumn technology 159
Nishimura, R., see Kominami, G. 216
Odham, G., see Sonesson, A. 11
Ogino, S., see Yanagi, H. 178
Oka, K., see Hirano, T. 371
O'Reilly, T.
—, Mac Mathúna, P., Keeling, P.W.N. and Feely, J.
Comparison of the spectrophotometric and high-performance liquid chromatographic analysis of Indocyanine Green in estimating systemic clearance in patients with chronic liver disease 190
Paanakker, J.E.
— and Van Hal, H.J.M.
Capillary gas chromatographic assay for the routine monitoring of the antidepressant mepirzepine in human plasma 203
Pachla, L.A., see Wright, D.S. 223
Palmerini, C.A.
—, Cantelmi, M.G., Minelli, A., Fini, C., Zampino, M. and Floridi, A.
Determination of plasma serotonin by high-performance liquid chromatography with pre-column sample enrichment and fluorimetric detection 378
Pauly, F., see Menu, P. 397
Persson, B.-A., see Balmér, K. 357
Petrarulo, M., see Mentasti, E. 253
Pinedo, H.M., see Maessen, P.A. 339
Porter, W.R., see Vanscoik, K.G. 183
Puglisi, C.V.
—, Ferrara, F.J. and Bekersky, I.
Determination of the agonist–antagonist anxiolytic agent 6-(2-chlorophenyl)-4-hydroxy-4*H*-imidazo[1,5-*a*][1,4]benzodiazepine-3-carboxamide and its benzophenone carboxylic acid metabolite in plasma by high-performance liquid chromatography 347
Ramzan, I.M.
Determination of the neuromuscular blocking drug gallamine in rat serum using high-performance liquid chromatography 428
Reams, G.P., see Souther, M.E. 27
Rejaibi, R., see Schöneshöfer, M. 434
Ress, K.
—, Liebich, H.M., Krämer, B., Ickrath, O., Risler, T. and Seipel, L.
Determination of amiodarone and its metabolite N-desethylamiodarone in serum by high-performance liquid chromatography – comparison of different extraction procedures 465
Richmond, R.
—, Turner, N.C., Maltby, N., Heavey, D., Vial, J., Dollery, C.T. and Taylor, G.W.
Single-step procedure for the extraction and purification of leukotrienes B_4, C_4 and D_4 241
Risler, T., see Ress, K. 465
Roberts, N.B., see Newton, C.J. 391
Ronfeld, R.A., see Fouda, H.G. 197
Rotatori, M., see Laganá, A. 135
Rothwell, C.M.
—, Hsieh, J.Y.-K. and Bayne, W.F.
High-performance liquid chromatographic determination of 4-[(3-(4-acetyl-3-hydroxy-2-propylphenoxy)propyl)sulfonyl]-γ-oxobenzenebutanoic acid and metabolites in human plasma 447
Sakai, J., see Togari, A. 41
Sakamoto, J., see Itami, T. 168
Sakoguchi, T., see Kobayashi, K. 439
Savigliano, M., see Mentasti, E. 253
Sayo, H.
—, Hatsumura, H., Hosokawa, M. and Michida, T.
Electrochemical immunoassay combined

with column liquid chromatography: determination of phenytoin in human serum 129
Schönberger, E., see MacCrehan, W.A. 65
Schöneshöfer, M.
—, Heilmann, P. and Rejaibi, R.
Automated column liquid chromatographic determination of polythiazide in human serum 434
Seipel, L., see Ress, K. 465
Shibati, K.
—, Kawada, T. and Iwai, K.
Microdetermination of N^1-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by high-performance liquid chromatography 173
Soest, R.E.J. van, see Nielen, M.W.F. 159
Soga, A., see Yanagi, H. 178
Sonesson, A.
—, Larsson, L. and Jimenez, J.
Use of pentafluorobenzyl and pentafluoropropionyl-pentafluorobenzyl esters of bacterial fatty acids for gas chromatographic analysis with electron-capture detection 366
Sonesson, A.
—, Larsson, L., Westerdahl, G. and Odham, G.
Determination of endotoxins by gas chromatography: evaluation of electron-capture and negative-ion chemical-ionization mass spectrometric detection of halogenated derivatives of β-hydroxymyristic acid 11
Souther, M.E.
—, Lumpkin, R.H., Kuo, K.C., Reams, G.P., Bauer, J.H. and Gehrke, C.W.
High-performance liquid chromatographic–radioimmunoassay method for the measurement of angiotensin II peptides in human plasma 27
Sugeno, K., see Kominami, G. 216
Sullivan, N.W.
— and Weiner, L.
Simple, rapid high-performance liquid chromatographic determination of bupivacaine in human plasma 236
Taniuchi, K., see Kobayashi, K. 439
Taylor, G.W., see Richmond, R. 241
Taylor, W.H., see Newton, C.J. 391
Thénot, J.J., see Dubruc, C. 208
Tieckelmann, H., see Ho, J. 269
Todo, S., see Tsunamoto, K. 414
Togari, A.
—, Sakai, J., Matsumoto, S. and Nagatsu, T.
Highly sensitive assay for alkaline and acid phosphatase activities by high-performance liquid chromatography with electrochemical detection 41
Tsunamoto, K.
—, Todo, S. and Imashuku, S.
Separation of prostaglandins and thromboxane by two-dimensional thin-layer chromatography 414
Turner, N.C., see Richmond, R. 241
Van der Vijgh, W.J.F., see Maessen, P.A. 339
Van Hal, H.J.M., see Paanakker, J.E. 203
Van Ingen, H.E., see Nielen, M.W.F. 159
Vanscoik, K.G.
—, Johnson, C.A. and Porter, W.R.
Sensitive and specific high-performance liquid chromatographic assay for 6-thiouric acid 183
Van Soest, R.E.J., see Nielen, M.W.F. 159
Vetter, B., see Benecke, R. 287
Vial, J., see Richmond, R. 241
Vigneron, C., see Menu, P. 397
Vijgh, W.J.F. van der, see Maessen, P.A. 339
Vree, T.B.
—, Baars, A.M. and De Grood, P.M.R.M.
High-performance liquid chromatographic determination and preliminary pharmacokinetics of propofol and its metabolites in human plasma and urine 458
Vries, J.X. de, see De Vries, J.X. 420
Wakasugi, K., see Hirano, T. 371
Walter-Sack, I., see De Vries, J.X. 420
Ward, A.E., see Bhamra, R.K. 229
Webster, H.K., see Krungkrai, J. 47
Webster, R.A., see Farrant, M. 385
Weidler, D.J., see Fouda, H.G. 197
Weiner, L., see Sullivan, N.W. 236
Westerdahl, G., see Sonesson, A. 11
Wright, D.S.
—, Pachla, L.A., Gibson, D.M. and Jordan, R.A.
Sensitive high-performance liquid chromatographic method for procaterol in human urine 223
Yamaguchi, T., see Kominami, G. 216
Yanagi, H.
—, Soga, A. and Ogino, S.
Purification of human urinary erythropoietin 178
Yunis, A.A., see Abou-Khalil, S. 111
Yuthavong, Y., see Krungkrai, J. 47
Zampino, M., see Palmerini, C.A. 378
Zeeuw, R.A. de, see Benecke, R. 287
Zhang, Y., see Balmér, K. 357
Zia-Gharib, F., see Farrant, M. 385
Zurlinden, E., see Conte, Jr., J.E. 452

Subject Index

4-[(3-(4-Acetyl-3-hydroxy-2-propylphenoxy)-propyl)sulfonyl]-γ-oxobenzenebutanoic acid
HPLC determination of 4-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propyl)sulfonyl]-γ-oxobenzenebutanoic acid and metabolites in human plasma 447
Acid phosphatase
Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 41
Aldehydes
HPLC determination of glyoxylic acid and other carbonyl compounds in urine 253
Alkaline phosphatase
Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 41
Amino acids
Automated pre-column derivatization with o-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase LC 385
Aminochloramphenicol
HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 111
δ-Aminolevulinic acid
Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 269
Amiodarone
Determination of amiodarone and its metabolite N-desethylamiodarone in serum by HPLC – comparison of different extraction procedures 465
Amphotericin B
HPLC determination of amphotericin B in human urine 439
Angiotensin II peptides
HPLC–radioimmunoassay method for the measurement of angiotensin II peptides in human plasma 27
Benzbromarone
Analysis of benzbromarone in human plasma and urine by HPLC and GC–MS 420
Benzodiazepines
Combined HPLC and enzyme immunoassay for active metabolites of a sleep inducer (rilmazafone), a ring-opened derivative of benzodiazepines, in human plasma 216
Benzoylecgonine
Determination of benzoylecgonine and cocaine in biologic fluids by automated GC 277
Betaxolol
Column LC measurement of betaxolol in plasma or serum 229
Bupivacaine
Simple, rapid HPLC determination of bupivacaine in human plasma 236
Cannabinoids
Direct injection of urine on a HPLC column-switching system for determination of Δ^9-tetrahydrocannabinol-11-oic acid with both UV and electrochemical detection 309
Captopril
Sensitive method for determination of captopril in biological fluids by GC–MS 79
Carbonyl compounds
HPLC determination of glyoxylic acid and other carbonyl compounds in urine 253
S-Carboxymethyl-L-cysteine
Determination of S-carboxymethyl-L-cysteine in plasma by HPLC with column switching following precolumn derivatization 208
Cefpiramide
Column LC determination of cefpiramide in human serum and urine 452
Chloramphenicol
HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 111
1-[2-(2-Chlorobenzoyl)phenyl]-4-(aminocarbonyl)-1*H*-imidazole-5-carboxylic acid
Determination of the agonist–antagonist anxiolytic agent 6-(2-chlorophenyl)-4-hydroxy-4*H*-imidazo[1,5-*a*][1,4]benzodiazepine-3-carboxamide and its benzophenone carboxylic acid metabolite in plasma by HPLC 347
6-(2-Chlorophenyl)-4-hydroxy-4*H*-imidazo-[1,5-*a*][1,4]benzodiazepine-3-carboxamide
Determination of the agonist–antagonist anxiolytic agent 6-(2-chlorophenyl)-4-hydroxy-4*H*-imidazo[1,5-*a*][1,4]benzodia-

zepine-3-carboxamide and its benzophenone carboxylic acid metabolite in plasma by HPLC 347

Cocaine
Determination of benzoylecgonine and cocaine in biologic fluids by automated GC 277

o,p'-DDD
Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 287

o,p'-DDE
Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 287

Deacetyldiltiazem
Assay of diltiazem and deacetyldiltiazem by capillary GC 89

Dehydrochloramphenicol
HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 111

7-Deoxydoxorubicin aglycone
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

7-Deoxydoxorubicinol aglycone
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

N-Desethylamiodarone
Determination of amiodarone and its metabolite N-desethylamiodarone in serum by HPLC – comparison of different extraction procedures 465

N-Desmethylzopiclone
Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by reversed-phase HPLC 151

Diamidines
Determination of fluorescent diamidines in plasma of experimental animals by HPLC 331

Dihydroergotamine
Determination of sub-nanogram amounts of dihydroergotamine in plasma and urine using LC and fluorimetric detection with off-line and on-line solid-phase drug enrichment 319

Diltiazem
Assay of diltiazem and deacetyldiltiazem by capillary GC 89

Doxorubicin aglycone
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

Doxorubicinol aglycone
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

Endotoxins
Determination of endotoxins by GC: evaluation of electron-capture and negative-ion chemical-ionization MS detection of halogenated derivatives of β-hydroxymyristic acid 11

Enzymes
Detection of uridine 5'-diphosphate glucose–collagen glucosyltransferase activity by HPLC and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 403

Enzymes
Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 41

Enzymes
HPLC assay for pteroylglutamate hydrolase 47

Enzymes
Separation of human pepsins in gastric juice by high-performance ion-exchange chromatography 391

4'-Epidoxorubicin
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

4'-Epidoxorubicin glucuronide
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

4'-Epidoxorubicinol
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

4'-Epidoxorubicinol glucuronide
Improved method for the determination of 4'-epidoxorubicin and seven metabolites in plasma by HPLC 339

Erythropoietin
Purification of human urinary erythropoietin 178

Ethynylestradiol
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 159

Ethynyl steroids
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 159
Fatty acids
Use of pentafluorobenzyl and pentafluoropropionyl-pentafluorobenzyl esters of bacterial fatty acids for GC analysis with electron-capture detection 366
α-Fetoprotein
Microheterogeneity of rat α-fetoprotein on a copper chelate column 168
Gallamine
Determination of the neuromuscular blocking drug gallamine in rat serum using HPLC 428
Glycoproteins
Microheterogeneity of rat α-fetoprotein on a copper chelate column 168
Glycoproteins
Purification of human urinary erythropoeitin 178
Glyoxylic acid
HPLC determination of glyoxylic acid and other carbonyl compounds in urine 253
Growth hormone
Fluorimetric determination of rat growth hormone in pituitary cell culture by HPLC with post-column derivatization 409
Haemoglobin pyridoxylation
Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate–haemoglobin species by HPLC resolution 397
α-Hydroxymetoprolol
Determination of metoprolol and two major metabolites in plasma and urine by column LC and fluorometric detection 357
β-Hydroxymyristic acid
Determination of endotoxins by GC: evaluation of electron-capture and negative-ion chemical-ionization MS detection of halogenated derivatives of β-hydroxymyristic acid 11
Indocyanine Green
Comparison of the spectrophotomeric and HPLC analysis of Indocyanine Green in estimating systemic clearance in patients with chronic liver disease 190
Iododeoxyuridine
Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 57
Keto acids
HPLC determination of glyoxylic acid and other carbonyl compounds in urine 253
Ketones
HPLC determination of glyoxylic acid and other carbonyl compounds in urine 253
Leukotrienes
Single-step procedure for the extraction and purification of leukotrienes B_4, C_4 and D_4 241
Lipid peroxides
LC determination of serum lipid peroxides as diphenylthiobarbituric acid-reactive substance to collagen disease patients – a preliminary study 371
Lipids
Single-step procedure for the extraction and purification of leukotrienes B_4, C_4 and D_4 241
Medroxalol
Application of column switching in HPLC analysis of medroxalol in plasma 143
Mepirzepine
Capillary GC assay for the routine monitoring of the antidepressant mepirzepine in human plasma 203
2-Mercaptopropionic acid
HPLC of 2-mercaptopropionylglycine and its metabolite 2-mercaptopropionic acid in plasma and urine after treatment with thiopronine 261
2-Mercaptopropionylglycine
HPLC of 2-mercaptopropionylglycine and its metabolite 2-mercaptopropionic acid in plasma and urine after treatment with thiopronine 261
N^1-Methyl-2-pyridone-5-carboxamide
Microdetermination of N^1-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 173
Metoprolol
Determination of metoprolol and two major metabolites in plasma and urine by column LC and fluorometric detection 357
Metoprolol acid
Determination of metoprolol and two major metabolites in plasma and urine by column LC and fluorometric detection 357
Mitotane
Rapid micromethod for the analysis of mitotane and its metabolite in plasma by GC with electron-capture detection 287

Neurotransmitters
Automated pre-column derivatization with *o*-phthalaldehyde for the determination of neurotransmitter amino acids using reversed-phase LC 385
Nicotinamide
Microdetermination of N^1-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 173
Nicotinic acid
Microdetermination of N^1-methyl-2-pyridone-5-carboxamide, a major metabolite of nicotinic acid and nicotinamide, in urine by HPLC 173
Nitrophenylaminopropanedione
HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 111
Nitrosochloramphenicol
HPLC determination of chloramphenicol and four analogues using reductive and oxidative electrochemical and UV detection 111
Norethindrone
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 159
Norfloxacin
HPLC determination of norfloxacin in human tissues and plasma with fluorescence detection 135
Norgestrel
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 159
Nucleobases
Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 57
N-Oxidezopiclone
Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by reversed-phase HPLC 151
Pepsins
Separation of human pepsins in gastric juice by high-performance ion-exchange chromatography 391
Peptides
HPLC–radioimmunoassay method for the measurement of angiotensin II peptides in human plasma 27
Peroxides
LC determination of serum lipid peroxides as diphenylthiobarbituric acid-reactive substance to collagen disease patients – a preliminary study 371
Phenytoin
Electrochemical immunoassay combined with column LC: determination of phenytoin in human serum 129
Phosphatases
Highly sensitive assay for alkaline and acid phosphatase activities by HPLC with electrochemical detection 41
Polythiazide
Automated column LC determination of polythiazide in human serum 434
Porphobilinogen
Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 269
Porphyrins
Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 269
Procaterol
Sensitive HPLC method for procaterol in human urine 223
Profolol
HPLC determination and preliminary pharmacokinetics of propofol and its metabolites in human plasma and urine 458
Propafenone
Determination of propafenone in biological fluids by fused-silica capillary GC using electron-capture detection 295
Prostaglandins
Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC–negative-ion chemical-ionization MS 1
Prostaglandins
Separation of prostaglandins and thromboxane by two-dimensional TLC 414
Prostanoids
Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC–negative-ion chemical-ionization MS 1
Pteroylpolyglutamate hydrolase
HPLC assay for pteroylglutamate hydrolase 47

Pyridoxal phosphate–haemoglobin
Haemoglobin pyridoxylation: optimization and observation of an additional new pyridoxal phosphate–haemoglobin species by HPLC resolution 397
Retinol
Reversed-phase HPLC separation and electrochemical detection of retinol and its isomers 65
Rilmazafone
Combined HPLC and enzyme immunoassay for active metabolites of a sleep inducer (rilmazafone), a ring-opened derivative of benzodiazepines, in human plasma 216
Serotonin
Determination of plasma serotonin by HPLC with pre-column sample enrichment and fluorimetric detection 378
Sertraline
GC–MS analysis and preliminary human pharmacokinetics of setraline, a new antidepressant drug 197
Steroids
Selective on-line trace enrichment for the determination of ethynyl steroids in urine by LC with precolumn technology 159
Δ^9-Tetrahydrocannabinol-11-oic acid
Direct injection of urine on a HPLC column-switching system for determination of Δ^9-tetrahydrocannabinol-11-oic acid with both UV and electrochemical detection 309
Theophylline
Ion-pair HPLC procedure for the quantitative analysis of theophylline in serum samples 99
6-Thiouric acid
Sensitive and specific HPLC assay for 6-thiouric acid 183
Thromboxane
Separation of prostaglandins and thromboxane by two-dimensional TLC 414
Thromboxane A_2
Purification and quantitative analysis of urinary prostanoids in human and in rat by packed and capillary GC–negative-ion chemical-ionization MS 1
Thymidine
Technique for detection of DNA nucleobases by reversed-phase HPLC optimized for quantitative determination of thymidine substitution by iododeoxyuridine 57
Uridine 5′-diphosphate glucose–collagen glucosyltransferase
Detection of uridine 5′-diphosphate glucose–collagen glucosyltransferase activity by HPLC and possible participation of a novel intermediate as a glucose donor in collagen biosynthesis 403
Vancomycin
HPLC analysis of vancomycin in plasma, bone, atrial appendage tissue and pericardial fluid 121
Vitamin A
Reversed-phase HPLC separation and electrochemical detection of retinol and its isomers 65
Xamoterol
Determination of xamoterol in human plasma by HPLC with electrochemical detection 233
Xanthines
Ion-pair HPLC procedure for the quantitative analysis of theophylline in serum samples 99
Zinc protoporphyrin
Quantitative determination of porphyrins, their precursors and zinc protoporphyrin in whole blood and dried blood by HPLC with fluorimetric detection 269
Zopiclone
Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by reversed-phase HPLC 151